

Orthospectra of Geodesic Laminations and Dilogarithm Identities on Moduli Space

Martin Bridgeman

March 4, 2009

Abstract

Given a measured lamination λ on a finite area hyperbolic surface we consider a natural measure M_λ on the real line obtained by taking the push-forward of the volume measure of the unit tangent bundle of the surface under an intersection function associated with the lamination. We show that the measure M_λ gives summation identities for the Rogers dilogarithm function on the moduli space of a surface.

1 Introduction

Let S be a closed hyperbolic surface and λ a geodesic lamination on S . We let Ω be the volume measure on the unit tangent bundle $T_1(S)$. We let $\alpha(v)$ be the longest geodesic arc containing v as a tangent vector and which does not intersect λ transversely in its interior. Generically $\alpha(v)$ will be a geodesic arc with endpoints on λ .

We define the function $L : T_1(S) \rightarrow \overline{\mathbf{R}}$ by letting $L(v) = \text{Length}(\alpha(v))$. We note that $L(v)$ is measurable but can be infinite. We define measure M_λ on the real line by $M_\lambda = L_*\Omega$. Then M_λ is a measure describing the distribution of the lengths of $\alpha(v)$.

We cut S along λ to obtain a surface with boundary denoted S_l . A λ -*cusp* of S is an ideal vertex of a component of S_l . We let N_λ be the number of λ -cusps of S . We denote by $\{\alpha_i\}$ the geodesic arcs in S_l which have endpoints perpendicular to $\partial S_l \subseteq \lambda$ and denote the length of α_i by l_i . We note that if a component of S_l is an ideal k -gon then there are a finite number of geodesics α_i in this component. Otherwise there are an infinite number. We call the set $\{l_i\}$ (with multiplicities) the λ -*orthospectrum*. By doubling $S - \lambda$ we see that the λ -orthospectrum corresponds to a subset of the closed geodesics of a finite area surface and therefore is a countable set.

We prove the following length spectrum identity

$$\sum_i \mathcal{L} \left(\frac{1}{\cosh^2 \frac{l_i}{2}} \right) = \frac{\pi^2}{12} (6|\chi(S)| - N_\lambda) \quad (1)$$

where $\mathcal{L}$ is a Rogers dilogarithm function (described below).

2 Dilogarithms and Polylogarithms

The k^{th} polylogarithm function Li_k is defined by the Taylor series

$$\text{Li}_k(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^k}$$

for $|z| < 1$ and by analytic continuation to C . In particular

$$\text{Li}_0(z) = \frac{1}{1-z} \quad \text{Li}_1(z) = -\log(1-z).$$

Also

$$\text{Li}'_k(z) = \frac{\text{Li}_{k-1}(z)}{z} \quad \text{giving} \quad \text{Li}_k(z) = \int_0^z \frac{\text{Li}_{k-1}(z)}{z} dz.$$

Also the functions Li_k are related to the Riemann ζ function by $\text{Li}_k(1) = \zeta(k)$.

The dilogarithm function is the function $\text{Li}_2(z)$ and is given by

$$\text{Li}_2(z) = - \int_0^z \frac{\log(1-z)}{z} dz.$$

Below is a brief description of some properties of the dilogarithm function. They can all be found in 1991 survey "Structural Properties of Polylogarithms" by L. Lewin (see [3]). From the power series representation, it is easy to see that the dilogarithm function satisfies the functional equation

$$\text{Li}_2(z) + \text{Li}_2(-z) = \frac{1}{2}\text{Li}_2(z^2).$$

Other functional relations of the dilogarithm can be best described by normalizing the dilogarithm function. The (extended) Rogers $\mathcal{L}$ -function (see [5]) is defined by

$$\mathcal{L}(x) = \text{Li}_2(x) + \frac{1}{2} \log|x| \log(1-x) \quad x \leq 1.$$

In terms of the Rogers $\mathcal{L}$ -function, Euler's reflection relations for the dilogarithm are

$$\begin{aligned} \mathcal{L}(x) + \mathcal{L}(1-x) &= \mathcal{L}(1) = \frac{\pi^2}{6} & 0 \leq x \leq 1 \\ \mathcal{L}(-x) + \mathcal{L}(-x^{-1}) &= 2\mathcal{L}(-1) = -\frac{\pi^2}{6} & x > 0 \end{aligned} \quad (2)$$

Also in terms of $\mathcal{L}$, Landen's identity is

$$\mathcal{L}\left(\frac{-x}{1-x}\right) = -\mathcal{L}(x) \quad 0 < x < 1 \quad (3)$$

and Abel's functional equation is

$$\mathcal{L}(x) + \mathcal{L}(y) = \mathcal{L}(xy) + \mathcal{L}\left(\frac{x(1-y)}{1-xy}\right) + \mathcal{L}\left(\frac{y(1-x)}{1-xy}\right). \quad (4)$$

Also a closed form for $\mathcal{L}(x)$ is known for certain values of x including

$$\mathcal{L}(1) = \frac{\pi^2}{6} \quad \mathcal{L}\left(\frac{1}{2}\right) = \frac{\pi^2}{12} \quad (\text{Euler}) \quad \mathcal{L}(\phi^{-1}) = \frac{\pi^2}{10} \quad \mathcal{L}(1-\phi^{-1}) = \frac{\pi^2}{15} \quad (\text{Landen})$$

where ϕ is the golden ratio.

3 Statement of Results

The main result of the paper is the following;

Main Theorem *There exists a function $\rho : \mathbf{R}^2 \rightarrow \mathbf{R}$ such that infinitesimally*

$$dM_\lambda = \left(\frac{4N_\lambda x^2}{\sinh^2 x} + \sum_i \rho(l_i, x) \right) dx$$

where N_λ is the number of λ -cusps of S . Furthermore the total mass of the measure $\rho(l, x)dx$ on the real line is given by

$$F(l) = \int_0^\infty \rho(l, x) dx = 8\mathcal{L} \left(\frac{1}{\cosh^2 \frac{l_i}{2}} \right)$$

In particular the measure M_λ depends only on the λ -orthospectrum.

4 Length Spectrum Identity

As $M_\lambda = L_*\Omega$, M_λ has total mass equal to the volume of $T_1(S)$. Therefore $M_\lambda(\mathbf{R}) = \Omega(T_1(S)) = 4\pi^2|\chi(S)|$. Summing up the masses of measures in the Main Theorem we immediately obtain the following.

Length Spectrum Identity Theorem *Let λ be a geodesic lamination on a finite area hyperbolic surface S . Then the λ -orthospectrum satisfies the following*

$$\sum_i \mathcal{L} \left(\frac{1}{\cosh^2 \frac{l_i}{2}} \right) = \frac{\pi^2(6|\chi(S)| - N_\lambda)}{12}$$

or equivalently

$$\sum_i \mathcal{L} \left(-\frac{1}{\sinh^2 \frac{l_i}{2}} \right) = \frac{\pi^2(6\chi(S) + N_\lambda)}{12}$$

By Landen's identity (see equation 3) we have

$$\mathcal{L} \left(\frac{1}{\cosh^2(\frac{l}{2})} \right) = -\mathcal{L} \left(-\frac{1}{\sinh^2(\frac{l}{2})} \right).$$

Thus we can see that the second form of the Length Spectrum Identity corresponds to the first via Landen's identity.

5 Length Spectrum Identity on Moduli Space

We note that if S is a connected hyperbolic surface of finite area with non-empty geodesic boundary, letting $\lambda = \partial S$ then the Length Spectrum Identity gives a summation identity on the Moduli space $Mod(S)$ of S . In this case the Euler characteristic $\chi(S)$ can be a fraction and is defined such that $2\pi\chi(S)$ is the negative of the area of S . This relation is an infinite relation except in the case when S is an ideal polygon. In this case we will show that these finite identities include the classical dilogarithm identities described above.

5.1 Classical Identities and the Moduli space of ideal polygons

For S an ideal n -gon, the Length Spectrum Identity is a finite summation relation. We will show that the associated relations give an infinite list of finite relations including the classical identities stated in the previous section.

If $\{l_i\}$ is a λ -orthospectrum, we will define two parameterizations by letting

$$a_i = -\frac{1}{\sinh^2 \frac{l_i}{2}} \quad b_i = \frac{1}{\cosh^2 \frac{l_i}{2}}.$$

We now consider the Poincaré disk model and let $x_i, i = 1, \dots, n$ be the vertices in anticlockwise cyclic ordering around the circle. Let s_i be the side $x_i x_{i+1}$. Let l_{ij} be the length of the diagonal between s_i and s_j for $|i - j| \geq 2$. We define the cross-ratio by

$$[z_1, z_2, z_3, z_4] = \frac{(z_1 - z_2)(z_4 - z_3)}{(z_1 - z_3)(z_4 - z_2)}.$$

As the cross ratio is invariant under Möbius transformations, we map the quadruple $(x_i, x_{i+1}, x_j, x_{j+1})$ to $(-1, 1, e^{l_{ij}}, -e^{l_{ij}})$. Then

$$[x_i, x_{i+1}, x_j, x_{j+1}] = [-1, 1, e^{l_{ij}}, -e^{l_{ij}}] = -\frac{(-1-1)(-e^{l_{ij}}-e^{l_{ij}})}{(-1-e^{l_{ij}})(-e^{l_{ij}}-1)} = \frac{4e^{l_{ij}}}{(e^{l_{ij}}+1)^2} = \frac{1}{\cosh^2(\frac{l_{ij}}{2})}$$

As S has area $(n-2)\pi$ and n cusps, $\chi(S) = (n-2)/2$ and $N_\lambda = n$. Thus the Length Spectrum identity becomes

$$\sum_{i,j} \mathcal{L}([x_i, x_{i+1}, x_j, x_{j+1}]) = \frac{(n-3)\pi^2}{6} \quad (5)$$

where the sum is over all ordered pairs i, j such that the sides s_i, s_j are disjoint (at infinity). In terms of dilogarithms we get

$$\sum_{i,j} \text{Li}_2([x_i, x_{i+1}, x_j, x_{j+1}]) = \frac{(n-3)\pi^2}{6} - \frac{1}{2} \sum_{i,j} \log(1 - [x_i, x_{i+1}, x_j, x_{j+1}]) \log([x_i, x_{i+1}, x_j, x_{j+1}]) \quad (6)$$

5.2 Some Cases

Quadrilateral: The ideal quadrilateral has 4 cusps and two ortholengths l_1, l_2 . By elementary hyperbolic geometry we have $\sinh(l_1/2) \cdot \sinh(l_2/2) = 1$. Therefore $a_1 \cdot a_2 = 1$ and letting $a = a_1$ the Length Spectrum identity is equivalent to the the classical reflection identity of Euler.

$$\mathcal{L}(a) + \mathcal{L}(a^{-1}) = -\frac{\pi^2}{6}. \quad (7)$$

Also we have

$$b_2 = \frac{1}{\cosh^2(l_2/2)} = \frac{1}{1 + \sinh^2(l_2/2)} = \frac{1}{1 + \frac{1}{\sinh^2(l_1/2)}} = \frac{\sinh^2(l_1/2)}{\cosh^2(l_1/2)} = 1 - \frac{1}{\cosh^2(l_1/2)} = 1 - b_1$$

Thus letting $b = b_1$, the Length Spectrum identity is equivalent to the Euler reflection identity

$$\mathcal{L}(b) + \mathcal{L}(1 - b) = \frac{\pi^2}{6}. \quad (8)$$

Pentagon and Abel's Identity: If we choose a general ideal pentagon then there are 5 diagonals and therefore 5 parameters a_i . We send three of the vertices to $0, 1, \infty$ and the other two to u, v with $0 < u < v < 1$. Then the cross ratios in terms of u, v are

$$u, \quad 1 - v, \quad \frac{v - u}{v}, \quad \frac{v - u}{1 - u}, \quad \frac{u(1 - v)}{v(1 - u)}$$

Putting into the equation we obtain the following equation.

$$\mathcal{L}(u) + \mathcal{L}(1 - v) + \mathcal{L}\left(\frac{v - u}{v}\right) + \mathcal{L}\left(\frac{v - u}{1 - u}\right) + \mathcal{L}\left(\frac{u(1 - v)}{v(1 - u)}\right) = \frac{\pi^2}{3}. \quad (9)$$

Letting $x = u/v, y = v$, then we get

$$\mathcal{L}(xy) + \mathcal{L}(1 - y) + \mathcal{L}(1 - x) + \mathcal{L}\left(\frac{y(1 - x)}{1 - xy}\right) + \mathcal{L}\left(\frac{x(1 - y)}{1 - xy}\right) = \frac{\pi^2}{3}. \quad (10)$$

Now by applying Euler's reflection identities for x, y , we obtain Abel's identity for the Rogers $\mathcal{L}$ -function.

$$\mathcal{L}(x) + \mathcal{L}(y) = \mathcal{L}(xy) + \mathcal{L}\left(\frac{y(1 - x)}{1 - xy}\right) + \mathcal{L}\left(\frac{x(1 - y)}{1 - xy}\right). \quad (11)$$

General equation: We obtain similar finite identities in the general ideal n -gon case. In general we note that equation 5 will have $(n - 3)$ independent variables and will be given by the summation of evaluating $\mathcal{L}$ on $\frac{n(n-3)}{2}$ rational functions in the $(n - 3)$ variables.

5.3 Regular Ideal n-gon relation

We now consider the dilogarithm equation for the specific case of a regular ideal n-gon. In this case the cross ratios can be calculated and the dilogarithm formulas for specific values of the dilogarithm function.

We consider a regular ideal n -gon in with center 0 in the Poincaré disk model and vertices at $v_k = u^k, k = 0, \dots, n-1$ for $u = e^{\frac{2\pi i}{n}}$. Then equation 5 can be thought of as an equation on the roots of the polynomial $z^n = 1$. We have

$$[v_0, v_1, v_r, v_{r+1}] = -\frac{(1-u)(u^{r+1}-u^r)}{(1-u^r)(u^{r+1}-u)} = \frac{u^r(u-1)^2}{u \cdot (u^r-1)^2} = \frac{\sin^2(\frac{\pi}{n})}{\sin^2(\frac{r\pi}{n})}$$

For $r < n/2$ there are exactly n distinct perpendiculars between sides separated by r sides and for $r = n/2$ there are $n/2$ such sides. To take care of the even and odd case simultaneously we let e_n be 1 if n is even and 0 if n is odd. Therefore we have

$$\sum_{r=2}^{\lceil n/2 \rceil - 1} n \cdot \mathcal{L} \left(\frac{\sin^2(\frac{\pi}{n})}{\sin^2(\frac{r\pi}{n})} \right) + e_n \frac{n}{2} \cdot \mathcal{L} \left(\sin^2 \left(\frac{\pi}{n} \right) \right) = \frac{(n-3)\pi^2}{6} \quad (12)$$

where $\lceil x \rceil$ is the least integer greater than or equal to x . Dividing by n we get

$$\sum_{r=2}^{\lceil n/2 \rceil - 1} \mathcal{L} \left(\frac{\sin^2(\frac{\pi}{n})}{\sin^2(\frac{r\pi}{n})} \right) + \frac{e_n}{2} \cdot \mathcal{L} \left(\sin^2 \frac{\pi}{n} \right) = \frac{(n-3)\pi^2}{6n}$$

Limiting case: We let n go to infinity and obtain the equation

$$\lim_{n \rightarrow \infty} \left(\sum_{r=2}^{\lceil n/2 \rceil - 1} \mathcal{L} \left(\frac{\sin^2(\frac{\pi}{n})}{\sin^2(\frac{r\pi}{n})} \right) + \frac{e_n}{2} \cdot \mathcal{L} \left(\sin^2 \frac{\pi}{n} \right) \right) = \lim_{n \rightarrow \infty} \frac{(n-3)\pi^2}{6n} = \frac{\pi^2}{6}$$

This gives a Rogers $\mathcal{L}$ -function series relation due to Lewin (see p. 298 of [3])

$$\sum_{r=2}^{\infty} \mathcal{L} \left(\frac{1}{r^2} \right) = \frac{\pi^2}{6}$$

Regular ideal quadrilateral: This case is trivial $a_1 = a_2 = -1$, $b_1 = b_2 = 1/2$ and equations 7,8 give the classical evaluations

$$\mathcal{L}(-1) = -\frac{\pi^2}{12} \quad \text{and} \quad \mathcal{L}\left(\frac{1}{2}\right) = \frac{\pi^2}{12}.$$

Regular ideal pentagon, Golden Mean: For the regular ideal pentagon, the orthospectrum consists of 5 geodesics each of the same length l . Using the formula above for $n = 5, r = 2$ we obtain that l satisfies

$$\cosh^2 \left(\frac{l}{2} \right) = \frac{2}{\sqrt{5} + 3} = \phi^2$$

where ϕ is the golden mean. Therefore as $\phi^2 = \phi + 1$

$$\sinh^2\left(\frac{l}{2}\right) = \phi^2 - 1 = \phi$$

and we have $a = -\phi^{-1}$. Thus the Length Spectrum Identity gives the classical relations of Landen

$$\mathcal{L}(-\phi^{-1}) = -\frac{\pi^2}{15} \quad \mathcal{L}(\phi^{-2}) = \frac{\pi^2}{15}.$$

Applying the quadrilateral relations 7, 8 we also get

$$\mathcal{L}(-\phi) = -\frac{\pi^2}{6} - \mathcal{L}(-\phi^{-1}) = -\frac{\pi^2}{10} \quad .$$

Regular ideal Hexagon: For a regular ideal hexagon, there are 9 elements of the orthospectrum, with the 6 being perpendicular to sides one apart and three being perpendicular to opposite sides. Putting $n = 6$ into equation 12 above then gives

$$6\mathcal{L}\left(\frac{1}{3}\right) + 3\mathcal{L}\left(\frac{1}{4}\right) = \frac{\pi^2}{2}$$

Before we prove the main theorem, we first consider the geometry of ideal quadrilaterals in the hyperbolic plane.

6 Intersections with ideal quadrilaterals

Given two disjoint geodesics g_1, g_2 with perpendicular distance l between them, let Q be the ideal quadrilateral with opposite sides g_1, g_2 . Then we can map Q by a Möbius transformation to the ideal quadrilateral Q_a in the upper half-plane with vertices $a, 0, 1, \infty \in \overline{\mathbf{R}}$ where $a < 0$. Similarly we can map Q to the ideal quadrilateral Q_b in the upper half-plane with vertices $0, b, 1, \infty \in \overline{\mathbf{R}}$ where $b > 0$. Using cross-ratios we have that

$$a = -\frac{1}{\sinh^2 \frac{l_a}{2}} \quad b = \frac{1}{\cosh^2 \frac{l_a}{2}} \quad (13)$$

The choice of normalization Q_a, Q_b leads to the equivalent forms of the Length Spectrum Identity. We choose normalization Q_a for our calculations.

If $x, y \in \mathbf{R}, x \neq y$, we let $g(x, y)$ be the geodesic in the upper half plane with end points x, y . Then for $(x, y) \in (a, 0) \times (1, \infty)$, the geodesic $g(x, y)$ intersects Q_a in a definite length denoted $L(x, y)$.




Figure 1: Length intersection

Lemma 1 *The map $L : (a, 0) \times (1, \infty) \rightarrow \mathbf{R}$ is given by the formula*

$$L(x, y) = \frac{1}{2} \ln \left(\frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right) = \frac{1}{2} \ln \frac{f(y)}{f(x)}.$$

where

$$f(x) = \frac{x(x-a)}{x-1}$$

Proof: Let T be the ideal triangle with vertices $0, 1, \infty$. Let $l_1 : (-\infty, 0) \times (0, 1) \rightarrow \mathbf{R}$ and $l_2 : (-\infty, 0) \times (1, \infty) \rightarrow \mathbf{R}$ be given by letting $l_1(x, y)$ be the length of the intersection of $g(x, y)$ with T and $l_2(x, y)$ be the length of the intersection of $g(x, y)$ with T . By a previous paper (see [1]) the functions l_i are given by

$$l_1(x, y) = \frac{1}{2} \ln \left(\frac{1-x}{1-y} \right) \quad l_2(x, y) = \frac{1}{2} \ln \left(\frac{y(x-1)}{x(y-1)} \right)$$

To calculate L , we split the quadrilateral Q_a by the vertical line at $x = 0$ into two ideal triangles T_1, T_2 where T_1 has vertices $0, 1, \infty$ and T_2 has vertices $a, 0, \infty$. Then $T_1 = T$ and $f_2(z) = z/a$ sends T_2 to T . Therefore

$$L(x, y) = l_2(x, y) + l_1(y/a, x/a)$$

Therefore

$$L(x, y) = \frac{1}{2} \ln \left(\frac{y(x-1)}{x(y-1)} \right) + \frac{1}{2} \ln \left(\frac{1-y/a}{1-x/a} \right) = \frac{1}{2} \ln \left(\frac{y(x-1)(a-y)}{x(y-1)(a-x)} \right)$$

■




Figure 2: Graph of function $f(x)$

We consider the rational function $f(x)$ defined above. Differentiating we have

$$f'(x) = \frac{(2x - a)(x - 1) - 1 \cdot (x^2 - ax)}{(x - 1)^2} = \frac{x^2 - 2x + a}{(x - 1)^2}$$

Therefore $f(x)$ has two critical points $1 \pm \sqrt{1 - a}$. We label the critical points $x_0 = 1 - \sqrt{1 - a}$ and $y_0 = 1 + \sqrt{1 - a}$ and note that x_0 is a maximum and y_0 a minimum.

7 Proof of Summation Identity

By definition

$$(L_*\Omega)(\phi) = \int_{T_1(S)} \phi(L(v)) d\Omega.$$

Let α, β be two arcs in S_l with endpoints on ∂S_l . Then we say $\alpha \sim \beta$ if they are homotopic relative to the boundary ∂S_l .

We define the sets $A_i = \{v \in T_1(S) | \alpha(v) \sim \alpha_i\}$. Also for each λ -cusp c we define $A_c = \{v \in T_1(S) | \alpha(v) \sim c\}$ where $\alpha(v) \sim c$ if $\alpha(v)$ can be homotoped (rel boundary) out the cusp c . Note that for $v \in A_i$ or $v \in A_c$, $L(v)$ is finite. Finally we define the set A_∞ to be all v not in any A_i or A_c . By definition, the sets A_i, A_c, A_∞ form a partition of $T_1(S)$. If we double S_l along its boundary, the geodesic arcs α_i correspond to a subset of the geodesics of the doubled surface. Therefore as the length spectrum of the doubled surface is countable, so is the collection of arcs α_i in S_l . Also, by ergodicity of geodesic flow on S (see [2]), the set A_∞ is a measure zero.

Therefore

$$(L_*\Omega)(\phi) = \sum_i \int_{A_i} \phi(L(v)) d\Omega + \sum_c \int_{A_c} \phi(L(v)) d\Omega.$$

We let

$$a_i = -\frac{1}{\sinh^2 \frac{l_i}{2}}$$

Then setting $Q_i = Q_{a_i}$, we have that Q_i is a quadrilateral with perpendicular of length l_i . We lift α_i to the upper half plane so that it is the perpendicular of length l_i in Q_i . We lift each λ -cusp c to the ideal vertex at infinity between the vertical geodesics $x = 0, x = 1$. Let T be the ideal triangle with vertices $0, 1, \infty \in \overline{\mathbf{R}}$.

If $v \in T_1(\mathbf{H}^2)$ in the upper half plane, we define $g(v)$ to be the geodesic with tangent vector v . We also denote the endpoints of $g(v)$ by $(x(v), y(v))$.

We lift the set A_i to the set $A'_i \subseteq T_1(Q_i)$. Then for $v \in A'_i$ the geodesic arc $\alpha'(v) = Q_i \cap g(v)$ is a lift of $\alpha(v)$. Similarly we lift A_c to the set $A'_c \subseteq T_1(T)$. Then for $v \in A'_c$ the geodesic arc $\alpha'(v) = T \cap g(v)$ is a lift of $\alpha(v)$. By abuse of notation we also let Ω be the volume measure on $T_1(\mathbf{H}^2)$. We parameterize $T_1(\mathbf{H}^2)$ by $(x, y, l) \in \overline{\mathbf{R}} \times \overline{\mathbf{R}} \times \mathbf{R}$ where (x, y, l) corresponds to the vector v such that $g(v)$ has ordered endpoints (x, y) and v has basepoint on $g(v)$ a distance l from the highest point of $g(v)$ in the upper half-plane. Then the volume form Ω can be written as (see [4])

$$d\Omega = \frac{2dx dy dl}{(x - y)^2}.$$

Therefore

$$\int_{A_c} \phi(L(v)) d\Omega = \int_{A'_c} \frac{2 \cdot \phi(L(v)) dx dy dl}{(x - y)^2}$$

We note that $L(v)$ only depends on the endpoints and therefore we can write $L(v) = L(x, y)$. If $v \in A'_c$ then either (x, y) or $(y, x) \in (-\infty, 0) \times (1, \infty)$. Integrating over l we have

$$\int_{A'_c} \frac{2 \cdot \phi(L(v)) dx dy dl}{(x - y)^2} = \int_{-\infty}^0 \int_1^\infty \frac{4 \cdot \phi(L(x, y)) L(x, y) dx dy}{(x - y)^2}.$$

By our previous paper [1]

$$\int_{-\infty}^0 \int_1^\infty \frac{4 \cdot \phi(L(x, y)) L(x, y) dx dy}{(x - y)^2} = \int_0^\infty \frac{4 \cdot \phi(L) \cdot L^2 dL}{\sinh^2 L}.$$

Thus as there are N_λ λ -cusps we have

$$\sum_c \int_{A_c} \phi(L(v)) d\Omega = N_\lambda \cdot \int_0^\infty \frac{4 \cdot \phi(L) \cdot L^2 dL}{\sinh^2 L} = M_\infty(\phi)$$

where M_∞ is the measure with infinitesimal

$$dM_\infty = \frac{4N_\lambda x^2 dx}{\sinh^2 x}.$$

Similarly we have by lifting A_i to A'_i that

$$\int_{A_i} \phi(L(v)) \, d\Omega = \int_{A'_i} \frac{2 \cdot \phi(L(v)) \, dx dy dl}{(x-y)^2}.$$

If $v \in A'_i$ then either (x, y) or $(y, x) \in (a_i, 0) \times (1, \infty)$. Integrating over l we have

$$\int_{A'_i} \frac{2 \cdot \phi(L(v)) \, dx dy dl}{(x-y)^2} = \int_{a_i}^0 \int_1^\infty \frac{4 \cdot \phi(L(x, y)) L(x, y) \, dx dy}{(x-y)^2}.$$

For $a < 0$ we define $M_a(\phi)$ to be the righthandside of the above equation. Then

$$M_a(\phi) = \int_a^0 \int_1^\infty \frac{4 \cdot \phi(L(x, y)) L(x, y) \, dx dy}{(x-y)^2}.$$

Then

$$M_\lambda = M_\infty + \sum_i M_{a_i}$$

As $M_\lambda = L_* \Omega$ it has total mass equal to the volume of $T_1(S)$ which is $4\pi^2 |\chi(S)|$. Therefore

$$\Omega(T_1(S)) = 4\pi^2 |\chi(S)| = M_\lambda(1) = M_\infty(1) + \sum_i M_{a_i}(1) \quad (14)$$

By an elementary calculation (see [1])

$$\int_0^\infty \frac{x^2 dx}{\sinh^2 x} = \frac{\pi^2}{6}.$$

Therefore

$$M_\infty(1) = \int_0^\infty \frac{4N_\lambda x^2 dx}{\sinh^2 x} = 4N_\lambda \int_0^\infty \frac{x^2 dx}{\sinh^2 x} = 4N_\lambda \cdot \frac{\pi^2}{6} = \frac{2N_\lambda \pi^2}{3}.$$

Using lemma 1 we substitute the formula for $L(x, y)$ to obtain

$$M_a(1) = \int_a^0 \int_1^\infty \frac{2 \cdot \log \left(\frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right) dx dy}{(x-y)^2}.$$

Then by equation 14 above we obtain

$$4\pi^2 |\chi(S)| = M_\infty(1) + \sum_i M_{a_i}(1) = \frac{2N_\lambda \pi^2}{3} + \sum_i F(l_i)$$

giving the summation identity

$$\sum_i F(l_i) = 4\pi^2 |\chi(S)| - \frac{2N_\lambda \pi^2}{3} = \frac{2\pi^2}{3} (6|\chi(S)| - N_\lambda) \quad (15)$$

■

8 Integral Calculation

In this section we find a formula for $F(l)$ by calculating an integral. We note that by the previous section, we already know that the function F satisfies the functional equation 15. We will make use of this to reduce F to the form we wish independent of using any classical dilogarithm relations.

Lemma 2 For $a < 0$

$$\int_a^0 \int_1^\infty \frac{\log \left| \frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right|}{(x-y)^2} dx dy = -4\mathcal{L}(a)$$

Proof: We let

$$G(a) = \int_a^0 \int_1^\infty \frac{\log \left| \frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right|}{(x-y)^2} dx dy$$

Integrating by parts we get

$$\int \frac{\log \left| \frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right|}{(x-y)^2} dx = -\frac{\log \left| \frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right|}{x-y} + \int \frac{1}{x-y} \left(\frac{1}{x-1} - \frac{1}{x} - \frac{1}{x-a} \right) dx.$$

Using

$$\int \frac{1}{(x-a)(x-b)} dx = \frac{1}{a-b} (\log |x-a| - \log |x-b|)$$

we get

$$\begin{aligned} \int \frac{\log \left| \frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right|}{(x-y)^2} dx &= -\frac{\log \left| \frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right|}{x-y} + \frac{1}{y-1} (\log |x-y| - \log |x-1|) + \\ &\quad -\frac{1}{y} (\log |x-y| - \log |x|) - \frac{1}{y-a} (\log |x-y| - \log |x-a|) \\ &= \frac{\log \left| \frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right|}{y-x} - \frac{\log |x-1|}{y-1} + \frac{\log |x|}{y} + \frac{\log |x-a|}{y-a} + \log |x-y| \left(\frac{1}{y-1} - \frac{1}{y} - \frac{1}{y-a} \right) \end{aligned}$$

We define

$$I(y) = \int_a^0 \frac{\log \left| \frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right|}{(x-y)^2} dx$$

To evaluate the improper integral $I(y)$ we gather the divergent terms to find their limits. Therefore

$$\begin{aligned} I(y) &= \lim_{x \rightarrow 0^-} \log |x| \left(\frac{1}{y} - \frac{1}{y-x} \right) - \lim_{x \rightarrow a^+} \log |x-a| \left(\frac{1}{y-a} - \frac{1}{y-x} \right) + \\ &\quad \frac{\log \left| \frac{y(y-a)}{a(y-1)} \right|}{y} + \frac{\log |a|}{y-a} + \log |y| \left(\frac{1}{y-1} - \frac{1}{y} - \frac{1}{y-a} \right) \\ &\quad - \frac{\log \left| \frac{y(y-a)(a-1)}{a(y-1)} \right|}{y-a} + \frac{\log |a-1|}{y-1} - \frac{\log |a|}{y} - \log |a-y| \left(\frac{1}{y-1} - \frac{1}{y} - \frac{1}{y-a} \right) \end{aligned}$$

By elementary calculus, both limits are zero. As $y > 1$ and $a < 0$, when we gather the remaining terms by common denominators and get

$$I(y) = \frac{-2\log(-a) + 2\log(y-a) - \log(y-1)}{y} + \frac{\log(1-a) + \log(y) - \log(y-a)}{y-1} + \frac{2\log(-a) - \log(1-a) - 2\log(y) + \log(y-1)}{y-a}$$

We now rewrite in the following form

$$I(y) = \left(\frac{\log(y)}{y-1} - \frac{\log(y-1)}{y} \right) + 2 \left(\frac{\log\left(\frac{y-a}{-a}\right)}{y} - \frac{\log\left(\frac{y}{-a}\right)}{y-a} \right) + \left(\frac{\log\left(\frac{y-1}{1-a}\right)}{y-a} - \frac{\log\left(\frac{y-a}{1-a}\right)}{y-1} \right). \quad (16)$$

Before we calculate the integral of $I(y)$ we note some properties of the dilogarithm. As the dilogarithm function Li_2 satisfies

$$\text{Li}_2(z) = - \int_0^z \frac{\log(1-t)}{t} dt$$

Then $\mathcal{L}$ has derivative

$$\mathcal{L}'(x) = \frac{d}{dx} \left(\text{Li}_2(x) + \frac{1}{2} \log|x| \log(1-x) \right) = -\frac{\log(1-x)}{x} + \frac{1}{2} \left(\frac{\log(1-x)}{x} - \frac{\log|x|}{1-x} \right) = -\frac{1}{2} \left(\frac{\log(1-x)}{x} + \frac{\log|x|}{1-x} \right)$$

Now if $a < b$, then on the interval $x > b$, we have $(b-x)/(b-a) < 0$. We define

$$J(x, a, b) = 2\mathcal{L} \left(\frac{b-x}{b-a} \right).$$

Then differentiating J we get

$$J'(x, a, b) = 2\mathcal{L}' \left(\frac{b-x}{b-a} \right) \cdot \frac{-1}{b-a} = \left(\frac{\log\left(\frac{x-a}{b-a}\right)}{b-x} + \frac{\log\left(\frac{x-b}{b-a}\right)}{x-a} \right) = \left(\frac{\log\left(\frac{x-b}{b-a}\right)}{x-a} - \frac{\log\left(\frac{x-a}{b-a}\right)}{x-b} \right)$$

We set

$$J(y) = -J(y, 0, 1) - 2J(y, a, 0) + J(y, a, 1).$$

Then from equation 16 we have that

$$J'(y) = -J'(y, 0, 1) - 2J'(y, a, 0) + J'(y, a, 1) = I(y).$$

Therefore we have an antiderivative for I and integrate to find G to get

$$G(a) = \int_1^\infty I(y) dy = J(y)|_1^\infty = \lim_{y \rightarrow \infty} J(y) - \lim_{y \rightarrow 1^+} J(y).$$

We let $\mathcal{L}_\infty$ be the limit if $\mathcal{L}(x)$ as x tends to $-\infty$. Therefore

$$\lim_{y \rightarrow 1^+} J(y) = -2\mathcal{L}(0) - 4\mathcal{L}(a^{-1}) + 2\mathcal{L}(0) = -4\mathcal{L}(a^{-1}) \quad \lim_{y \rightarrow \infty} J(y) = -4\mathcal{L}_\infty.$$

Thus

$$G(a) = -4\mathcal{L}_\infty + 4\mathcal{L}(a^{-1}) = -4(\mathcal{L}_\infty - \mathcal{L}(a^{-1}))$$

It follows immediately from Euler's reflection identity that $G(a) = -4\mathcal{L}(a)$ but for completeness we derive it independently. From the formula we have $G(0) = -4(\mathcal{L}_\infty - \mathcal{L}_\infty) = 0$. Also by equation 15, $G(a)$ must satisfy a summation identity

$$G(a) + G(a^{-1}) = G_\infty$$

where G_∞ is the limit of G as a tends to $-\infty$. Therefore

$$G(a) = G_\infty - G(a^{-1}) = (G_\infty + 4\mathcal{L}_\infty) - 4\mathcal{L}(a)$$

But as $G(0) = 0$ we have

$$G(a) = -4\mathcal{L}(a) \quad \text{and finally} \quad F(l) = -8\mathcal{L}\left(-\frac{1}{\sinh^2(l/2)}\right).$$

■

We note that by performing the integral over the quadrilateral Q_b where $b = 1/\cosh^2(l/2)$, the above can be repeated to show

$$F(l) = 8\mathcal{L}\left(\frac{1}{\cosh^2(l/2)}\right).$$

Equivalently we note it also follows from Landen's identity.

9 Volume interpretation of $\mathcal{L}$

Let g_1, g_2 be disjoint geodesics in $\mathbf{H}^2$ with perpendicular distance l and endpoints x_1, y_1 and x_2, y_2 respectively on $\mathbf{S}^1$. Given $v \in T_1(S)$ let g_v be the associated oriented geodesic with tangent v . Then we define the set

$$C(g_1, g_2) = \{v \in T_1(S) \mid g_v \cap g_1 \neq \emptyset, g_v \cap g_2 \neq \emptyset\}$$

Let $t = [x_1, y_1, x_2, y_2]$, then depending on the ordering of the points on the circle we have

$$t = [-1, 1, e^l, -e^l] = \frac{1}{\cosh^2(l/2)} \quad \text{or} \quad t = [-1, 1, -e^l, e^l] = -\frac{1}{\sinh^2(l/2)}.$$

It follows from the invariance of volume on $T_1(S)$, that the volume of $C(g_1, g_2)$ in $T_1(S)$ only depends on t . We therefore define $V(t) = \text{Volume}(S(g_1, g_2))$.

Then it follows from the main theorem that

$$\mathcal{L}(t) = \pm \frac{1}{8} V(t)$$

where the sign is given by the sign of t . Therefore we can interpret the Rogers $\mathcal{L}$ -function as a signed volume function on $T_1(S)$ for the sets $G(g_1, g_2)$.

10 Integral Formula for ρ

We let

$$L(x, y) = \frac{1}{2} \log \left(\frac{y(y-a)(x-1)}{x(x-a)(y-1)} \right) = \frac{1}{2} \log \left(\frac{f(y)}{f(x)} \right) \quad \text{for} \quad f(x) = \frac{x(x-a)}{x-1}.$$

Taking derivatives of the length function $L(x, y)$ we have

$$\frac{\partial L}{\partial x} = -\frac{f'(x)}{2f(x)} \quad \frac{\partial L}{\partial y} = \frac{f'(y)}{2f(y)}.$$

By the previous section, the function f has critical points x_0, y_0 . Furthermore on $(a, 0)$ the function $f(x)$ has global maximum at x_0 and on $(1, \infty)$, f has global minimum at y_0 . Therefore fixing x , the function $u : (1, \infty) \rightarrow \mathbf{R}$ given by $u(y) = L(x, y)$ is decreasing on $(1, y_0)$ and increasing on (y_0, ∞) . Therefore we make the change of variable $t = L(x, y), x = x$. Finding inverses for f we define the two function g_+, g_- by

$$g_{\pm}(x) = \frac{(a+x) \pm \sqrt{(a+x)^2 - 4x}}{2}.$$

Then solving $t = L(x, y)$ gives $f(y) = f(x)e^{2t}$. Therefore on $(1, y_0)$ we have $y = g_-(f(x)e^{2t})$ and on (y_0, ∞) we have $y = g_+(f(x)e^{2t})$. Therefore

$$M_a(\phi) = \int_a^0 \left(\int_1^{y_0} + \int_{y_0}^{\infty} \frac{4 \cdot \phi(L(x, y)) L(x, y) dy}{(x-y)^2} \right) dx.$$

and

$$\begin{aligned} \int_1^{y_0} \frac{4 \cdot \phi(L(x, y)) L(x, y) dy}{(x-y)^2} &= \int_{\infty}^{L(x, y_0)} \frac{4 \cdot \phi(t) t \cdot g'_-(f(x)e^{2t}) 2f(x)e^{2t} dt}{(x - g_-(f(x)e^{2t}))^2} \\ \int_{y_0}^{\infty} \frac{4 \cdot \phi(L(x, y)) L(x, y) dy}{(x-y)^2} &= \int_{L(x, y_0)}^{\infty} \frac{4 \cdot \phi(t) t \cdot g'_+(f(x)e^{2t}) 2f(x)e^{2t} dt}{(x - g_+(f(x)e^{2t}))^2} \end{aligned}$$

Therefore combining we have

$$M_a(\phi) = \int_a^0 \int_{L(x, y_0)}^{\infty} 8 \cdot \phi(t) \cdot t \cdot e^{2t} \cdot f(x) \left(\frac{g'_+(f(x)e^{2t})}{(x - g_+(f(x)e^{2t}))^2} - \frac{g'_-(f(x)e^{2t})}{(x - g_-(f(x)e^{2t}))^2} \right) dt dx$$

We switch the order of integration. The function $L(x, y_0)$ is minimum at x_0 with minimum value $l = L(x_0, y_0)$ being the length of the perpendicular (see figure 2). Thus we integrate t from l to infinity. The integral in the x direction is between the two x solutions of $t = L(x, y_0)$ which are solutions to $f(x) = f(y_0)e^{-2t}$. Thus we integrate x from $g_-(f(y_0)e^{-2t})$ to $g_+(f(y_0)e^{-2t})$ giving

$$M_a(\phi) = \int_{l_a}^{\infty} 8 \cdot \phi(t) \cdot t \cdot e^{2t} dt \left(\int_{g_-(f(y_0)e^{-2t})}^{g_+(f(y_0)e^{-2t})} \left(\frac{g'_+(f(x)e^{2t})}{(x - g_+(f(x)e^{2t}))^2} - \frac{g'_-(f(x)e^{2t})}{(x - g_-(f(x)e^{2t}))^2} \right) f(x) dx \right)$$

Therefore

$$M_a(\phi) = \int_0^{\infty} \phi(t) \cdot \rho(l, t) dt$$

where

$$\rho(l, t) = 8te^{2t} \chi_{[l, \infty)} \cdot \left(\int_{g_-(f(y_0)e^{-2t})}^{g_+(f(y_0)e^{-2t})} \left(\frac{g'_+(f(x)e^{2t})}{(x - g_+(f(x)e^{2t}))^2} - \frac{g'_-(f(x)e^{2t})}{(x - g_-(f(x)e^{2t}))^2} \right) f(x) dx \right)$$

and $f(x) = \frac{x(x-a)}{x-1}$ where $a = -\frac{1}{\sinh^2 l/2}$

Therefore

$$(L_*\Omega)(\phi) = \int_0^\infty \phi(x) \rho(x) dx$$

where

$$\rho(x) = \frac{4N_\lambda x^2}{\sinh^2 x} + \sum_i \rho(l_i, x)$$

■

11 Asymptotic behavior

In this section we study the asymptotic behavior of the function $\rho(l, t)$ for large t .

For functions of a single variable, we write $f(x) \simeq g(x)$ as x tends to x_0 if

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 1.$$

Furthermore for functions of more than one variable, we write $f(x, y) \simeq_x g(x, y)$ as x tends to x_0 if

$$\lim_{x \rightarrow x_0} \frac{f(x, y)}{g(x, y)} = 1.$$

Theorem 3 *The measure $\rho(l, t)dx$ on the real line satisfies*

$$\lim_{t \rightarrow \infty} \frac{\rho(l, t)}{16t^2 e^{-2t}} = r(l)$$

uniformly on compact subsets of $(0, \infty)$ where

$$r(l) = \frac{-2a^2 + 5a - 2}{a(1-a)} \quad \text{for} \quad a = -\frac{1}{\sinh^2(l/2)}$$

Proof: We now show $\lim_{t \rightarrow \infty} \rho(l, t) = r(l)$ converges uniformly on compact subsets of $(0, \infty)$. Let $I \subseteq (0, \infty)$ be a compact interval. Now let $l \in I$. As before we let $a = -1/\sinh^2(l/2)$ and define $f(x) = x(x-a)/(x-1)$ with inverses $g_\pm$ and critical values x_0, y_0 . Let

$$G(t, x) = 8te^{2t} \left(\frac{g'_+(f(x)e^{2t})}{(x - g_+(f(x)e^{2t}))^2} - \frac{g'_-(f(x)e^{2t})}{(x - g_-(f(x)e^{2t}))^2} \right) f(x)$$

Then for $t > l$ we have

$$\rho(l, t) = \int_{g_-(f(y_0)e^{-2t})}^{g_+(f(y_0)e^{-2t})} G(t, x) dx$$

For $C > 0$, we further define

$$\rho(C, l, t) = \int_{g_-(f(y_0)Ce^{-2t})}^{g_+(f(y_0)Ce^{-2t})} G(t, x) dx \quad (17)$$

On the interval $[a, 0]$ f has maximum at x_0 . Therefore $\rho(C, l, t)$ is defined for all t such that $f(y_0)Ce^{-2t} < f(x_0)$ or

$$t > K_0(C) = \frac{1}{2} \ln C + \frac{1}{2} \ln \left(\frac{f(y_0)}{f(x_0)} \right) = l + \frac{1}{2} \ln C$$

Considering $g_{\pm}(x)$ for large x we have

$$g_{\pm}(x) = \frac{(a+x) \pm \sqrt{(a+x)^2 - 4x}}{2} \simeq \frac{(a+x)}{2} \left(1 \pm \left(1 - \frac{2x}{(a+x)^2} \right) \right)$$

Therefore

$$g_-(x) \simeq \frac{(a+x)}{2} \left(1 - 1 + \frac{2x}{(a+x)^2} \right) = \frac{x}{a+x} \simeq 1 - \frac{a}{x}$$

and

$$g_+(x) \simeq \frac{(a+x)}{2} \left(1 + 1 - \frac{2x}{(a+x)^2} \right) = (a+x) - \frac{x}{a+x} \simeq (a-1) + x + \frac{a}{x}$$

Taking leading terms we have

$$g_-(x) \simeq 1 \quad g'_-(x) \simeq \frac{a}{x^2} \quad g_+(x) \simeq x \quad g'_+(x) \simeq 1 \quad (18)$$

We let $I_C = [g_-(f(y_0)Ce^{-2t}), g_+(f(y_0)Ce^{-2t})]$. Then for $x \in I_C$ we have $f(x)e^{2t} \geq C.f(y_0)$. Therefore for C sufficiently large we use the above approximations to approximate $G(t, x)$ on I_C . We substitute the approximations 18 into the formula for $G(t, x)$ to define

$$G_1(t, x) = 8te^{2t} \cdot \left(\frac{1}{(x - f(x)e^{2t})^2} - \frac{\frac{a}{(f(x)e^{2t})^2}}{(x-1)^2} \right) f(x)$$

Simplifying we have

$$G_1(t, x) = 8te^{-2t} \cdot \left(\frac{1}{(1 - \frac{x}{f(x)e^{2t}})^2} - \frac{a}{(x-1)^2} \right) \frac{1}{f(x)}.$$

Noting that $f(x)e^{2t} > Cf(y_0)$ on I_C , then for large C the quantity $\frac{x}{f(x)e^{2t}}$ is small and we obtain the approximation

$$G_2(t, x) = 8te^{-2t} \cdot \left(1 - \frac{a}{(x-1)^2} \right) \frac{1}{f(x)}.$$

Therefore given an $\epsilon > 0$ we can find a $K_1(\epsilon)$ such that

$$\frac{G(t, x)}{G_2(t, x)} \in [1 - \epsilon, 1 + \epsilon] \quad \text{for all } C > K_1(\epsilon), t > K_0(C), x \in I_C.$$

Therefore integrating

$$\frac{1}{\rho(C, l, t)} \left(8te^{-2t} \cdot \int_{g_-(f(y_0)Ce^{-2t})}^{g_+(f(y_0)Ce^{-2t})} \left(1 - \frac{a}{(x-1)^2} \right) \frac{1}{f(x)} dx \right) \in [1 - \epsilon, 1 + \epsilon]$$

for $C > K_1(\epsilon)$ and $t > K_0(C)$. We fix a $K > K_1(\epsilon)$ and define

$$\begin{aligned} \rho_K(l, t) &= 8te^{-2t} \cdot \left(\int_{g_-(f(y_0)Ke^{-2t})}^{g_+(f(y_0)Ke^{-2t})} \left(1 - \frac{a}{(x-1)^2} \right) \frac{1}{f(x)} dx \right) \\ &= 8te^{-2t} \cdot \left(\int_{g_-(f(y_0)Ke^{-2t})}^{g_+(f(y_0)Ke^{-2t})} \left(\frac{x-1}{x \cdot (x-a)} - \frac{a}{x(x-a)(x-1)} \right) dx \right) \end{aligned}$$

Integrating we have

$$\int \left(\frac{x-1}{x \cdot (x-a)} - \frac{a}{x(x-a)(x-1)} \right) dx = \left(\frac{1-a}{a} \ln|x| - \frac{a}{1-a} \ln|x-1| - \frac{a^2-3a+1}{a(1-a)} \ln|x-a| \right)$$

Therefore

$$\rho_K(l, t) = 8te^{-2t} \cdot \left(\frac{1-a}{a} \ln|x| - \frac{a}{1-a} \ln|x-1| - \frac{a^2-3a+1}{a(1-a)} \ln|x-a| \right) \Big|_{g_-(f(y_0)Ke^{-2t})}^{g_+(f(y_0)Ke^{-2t})}$$

For x small we have

$$g_{\pm}(x) = \frac{(a+x) \pm \sqrt{(a+x)^2 - 4x}}{2} \simeq \frac{(a+x)}{2} \left(1 \mp \left(1 - \frac{2x}{(a+x)^2} \right) \right)$$

Therefore

$$g_-(x) \simeq (a+x) - \frac{x}{a+x} \simeq a - \frac{(1-a)x}{a} \quad g_+(x) \simeq \frac{x}{a+x} \simeq \frac{x}{a}$$

Therefore

$$\rho_K(l, t) \simeq_t 8te^{-2t} \cdot \left(\frac{1-a}{a} \ln \left| \frac{Kf(y_0)e^{-2t}}{a^2} \right| - \frac{a}{1-a} \ln \left| \frac{1}{a-1} \right| - \frac{a^2-3a+1}{a(1-a)} \ln \left| \frac{a^2}{(1-a)f(y_0)Ke^{-2t}} \right| \right)$$

Taking limits as we have

$$\rho_K(l, t) \simeq_t (16t^2e^{-2t}) \cdot \left(-\frac{1-a}{a} - \frac{a^2-3a+1}{a(1-a)} \right) = (16t^2e^{-2t}) \cdot \frac{-2a^2+5a-2}{a(1-a)}$$

Therefore given $\epsilon > 0$ there exists $K_1(\epsilon) > 0$ such that for any $C > K_1(\epsilon)$ both

$$\liminf_{t \rightarrow \infty} \frac{\rho(C, l, t)}{16t^2e^{-2t}r(a)} \quad \text{and} \quad \limsup_{t \rightarrow \infty} \frac{\rho(C, l, t)}{16t^2e^{-2t}r(a)} \quad \text{are in } [1 - \epsilon, 1 + \epsilon]. \quad (19)$$

where

$$r(a) = \frac{-2a^2+5a-2}{a(1-a)}$$

We now define

$$\rho_-(C, l, t) = \int_{g_-(f(y_0)e^{-2t})}^{g_-(f(y_0)Ce^{-2t})} G(t, x) dx \quad \text{and} \quad \rho_+(C, l, t) = \int_{g_+(f(y_0)Ce^{-2t})}^{g_+(f(y_0)e^{-2t})} G(t, x) dx.$$

Then by definition

$$\rho(l, t) = \rho(C, l, t) + \rho_-(C, l, t) + \rho_+(C, l, t).$$

We now bound the functions $\rho_\pm(C, l, t)$. Let I_C^-, I_C^+ be the given intervals.

On the interval I , $g_\pm(f(x)e^{2t}) > 1$ and $x < 0$ so $(x - g_\pm(f(x)e^{2t}))^2 > 1$. Also as $g'_-(f(x)e^{2t}) < 0$ we have

$$\begin{aligned} |G(t, x)| &= (8t.e^{2t}). \left(\frac{g'_+(f(x)e^{2t})}{(x - g_+(f(x)e^{2t}))^2} - \frac{g'_-(f(x)e^{2t})}{(x - g_-(f(x)e^{2t}))^2} \right) f(x). \\ &\leq 8t.e^{2t}. (g'_+(f(x)e^{2t}) - g'_-(f(x)e^{2t})) f(x). \end{aligned}$$

The derivative of $g_\pm(x)$ is given by

$$g'_\pm(x) = \frac{1}{2} \pm \frac{1}{2} \frac{x + a - 2}{\sqrt{(a + x)^2 - 4x}}.$$

Therefore

$$g'_+(x) - g'_-(x) = \frac{x + a - 2}{\sqrt{(a + x)^2 - 4x}}.$$

As f has critical values $f(x_0)$ and $f(y_0)$ we have that

$$g'_+(x) - g'_-(x) = \frac{x + a - 2}{\sqrt{(x - f(x_0))(x - f(y_0))}}.$$

We note that on $I_C^\pm$ we have $f(y_0) < f(x)e^{2t} < Cf(y_0)$ then

$$\begin{aligned} g'_+(f(x)e^{2t}) - g'_-(f(x)e^{2t}) &\leq \frac{Cf(y_0) + a - 2}{\sqrt{(f(y_0) - f(x_0))(f(x)e^{2t} - f(y_0))}}. \\ &\leq \left(\frac{Cf(y_0) + a - 2}{\sqrt{f(y_0) - f(x_0)}} \right) \frac{e^{-t}}{\sqrt{f(x) - f(y_0)e^{-2t}}} \end{aligned}$$

The function $f(x) = x(x - a)/(x - 1)$ has maximum at x_0 on $(a, 0)$. Therefore for $b < f(x_0)$

$$f(x) - b = \frac{(x - g_-(b))(x - g_+(b))}{(x - 1)}$$

As $x \in (a, 0)$ we have

$$f(x) - b \geq (x - g_-(b))(g_+(b) - x)$$

Therefore

$$g'_+(f(x)e^{2t}) - g'_-(f(x)e^{2t}) \leq \left(\frac{Cf(y_0) + a - 2}{\sqrt{f(y_0) - f(x_0)}} \right) \frac{e^{-t}}{\sqrt{(x - g_-(f(y_0)e^{-2t}))(g_+(f(y_0)e^{-2t}) - x)}}$$

Now restricting to I_C^+ we have $x > g_+(f(y_0)Ce^{-2t})$. Therefore for $x \in I_C^+$,

$$g'_+(f(x)e^{2t}) - g'_-(f(x)e^{2t}) \leq \left(\frac{Cf(y_0) + a - 2}{\sqrt{(f(y_0) - f(x_0))(g_+(f(y_0)Ce^{-2t}) - g_-(f(y_0)e^{-2t}))}} \right) \frac{e^{-t}}{\sqrt{g_+(f(y_0)e^{-2t}) - x}}$$

Therefore we have

$$\rho_+(C, l, t) \leq \int_{I_C^+} |G(t, x)| dx \leq D(t) 8te^t \cdot \int_{I_C^+} \frac{f(x)}{\sqrt{g_+(f(y_0)e^{-2t}) - x}} dt$$

where $D(t)$ is the constant

$$D(t) = \left(\frac{Cf(y_0) + a - 2}{\sqrt{(f(y_0) - f(x_0))(g_+(f(y_0)Ce^{-2t}) - g_-(f(y_0)e^{-2t}))}} \right)$$

As $f(x) = x(x - a)/(x - 1)$ then, $0 < f(x) \leq ax$ on $(a, 0)$ we have

$$\rho_+(C, l, t) \leq \int_{I_C^+} |G(t, x)| dx \leq D(t) \cdot 8 \cdot a \cdot t \cdot e^t \cdot \int_{I_C^+} \frac{x}{\sqrt{g_+(f(y_0)e^{-2t}) - x}} dx$$

By integration we have

$$\int_a^b \frac{x}{\sqrt{b - a}} dx = \frac{2}{3} (2b + a) \sqrt{b - a}$$

Therefore

$$\rho_+(C, l, t) \leq 16 \cdot D(t) \cdot a \cdot t \cdot e^t \cdot (2g_+(f(y_0)e^{-2t}) + g_+(f(y_0)Ce^{-2t})) \sqrt{g_+(f(y_0)e^{-2t}) - g_+(f(y_0)Ce^{-2t})}.$$

Now for t large we have

$$\lim_{t \rightarrow \infty} D(t) = \left(\frac{Cf(y_0) + a - 2}{\sqrt{(f(y_0) - f(x_0)) \cdot |a|}} \right) = D.$$

We note for x small $g_+(x) \simeq x/a$. Therefore

$$\begin{aligned} \limsup_{t \rightarrow \infty} \left| \frac{\rho_+(C, l, t)}{t^2 e^{-2t}} \right| &\leq \limsup_{t \rightarrow \infty} \frac{16 \cdot D \cdot a \cdot t \cdot e^t \cdot \left(\frac{2f(y_0)e^{-2t} + f(y_0)Ce^{-2t}}{a} \right) \sqrt{\frac{f(y_0)e^{-2t} - f(y_0)Ce^{-2t}}{a}}}{t^2 e^{-2t}}. \\ \limsup_{t \rightarrow \infty} \left| \frac{\rho_+(C, l, t)}{t^2 e^{-2t}} \right| &\leq \limsup_{t \rightarrow \infty} \frac{16 \cdot D \cdot f(y_0)^{3/2} (C + 2) \sqrt{C - 1}}{t \cdot \sqrt{-a}} = 0. \end{aligned}$$

Thus

$$\lim_{t \rightarrow \infty} \frac{\rho_+(C, l, t)}{t^2 e^{-2t}} = 0.$$

Similarly for $\rho_-(C, l, t)$ we once again have that

$$\lim_{t \rightarrow \infty} \frac{\rho_-(C, l, t)}{t^2 e^{-2t}} = 0.$$

Therefore given $\epsilon > 0$ we can find $K(\epsilon)$ such that for $C > K(\epsilon)$ by equations 19

$$\limsup_{t \rightarrow \infty} \frac{\rho(l, t)}{16t^2 e^{-2t} r(a)} = \limsup_{t \rightarrow \infty} \left(\frac{\rho_-(C, l, t)}{16t^2 e^{-2t} r(a)} + \frac{\rho(C, l, t)}{16t^2 e^{-2t} r(a)} + \frac{\rho_+(C, l, t)}{16t^2 e^{-2t} r(a)} \right) = \limsup_{t \rightarrow \infty} \frac{\rho(C, l, t)}{16t^2 e^{-2t} r(a)} \in [1 - \epsilon, 1 + \epsilon]$$

As ϵ is arbitrary we have

$$\limsup_{t \rightarrow \infty} \frac{\rho(l, t)}{16t^2 e^{-2t} r(a)} = 1$$

Similarly

$$\liminf_{t \rightarrow \infty} \frac{\rho(l, t)}{16t^2 e^{-2t} r(a)} = 1$$

■

References

- [1] Martin Bridgeman, David Dumas, Distribution of intersection lengths of a random geodesic with a geodesic lamination. *Ergodic Theory and Dynamical Systems*, 27(4), 2007
- [2] Eberhard Hopf. Statistik der geodätischen Linien in Mannigfaltigkeiten negativer Krümmung. *Ber. Verh. Sächs. Akad. Wiss. Leipzig*, 91:261–304, 1939.
- [3] L. Lewin, (Ed.). Structural Properties of Polylogarithms, *Mathematical Surveys and Monographs*, AMS, Providence, RI, 1991.
- [4] Peter J. Nicholls. *The Ergodic Theory of Discrete Groups*, volume 143 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1989.
- [5] L.J. Rogers. On Function Sum Theorems Connected with the Series $\sum_1^\infty \frac{x^n}{n^2}$ *Proc. London Math. Soc.* 4, 169-189, 1907